

Radiative constraints on the habitability of exoplanets Gliese 581c and Gliese 581d

Y. Hu^{1,2} and F. Ding^{1,2}

¹ Laboratory for Climate and Ocean-Atmosphere Studies, Peking University, 100871 Beijing, PR China
e-mail: yyhu@pku.edu.cn

² Dept. of Atmospheric and Oceanic Sciences, School of Physics, Peking University, 100871 Beijing, PR China

Received 28 April 2010 / Accepted 18 November 2010

ABSTRACT

Aims. The M-type star Gliese 581 is likely to have two super-Earth planets, i.e., Gl 581c and Gl 581d. The present study is to investigate their habitability constrained by radiative properties of their atmospheres and the threshold of carbon-dioxide (CO₂), assuming that the two exoplanets are terrestrial, and that they have similar outgassing processes to those of the terrestrial planets in our own solar system.

Methods. Radiative-convective atmospheric models are used. Different values of CO₂ concentrations and water-vapor mixing ratios are tested.

Results. Our simulation results suggest that Gl 581d is probably a habitable planet. However, at least 6.7 bars of CO₂ are required to raise its surface temperature (T_s) above the freezing point of water. In contrast, Gl 581c might have experienced runaway greenhouse, like Venus, because of its too high surface temperature and the lack of an effective cold trap for water vapor. We compare our results with other independent studies.

Key words. planetary systems – radiative transfer – planets and satellites: atmospheres – stars: individual: Gl 581

1. Introduction

Two super-Earth extra-solar planets were detected orbiting M-dwarf Gliese (Gl) 581 (Udry et al. 2007), called Gl 581c and Gl 581d. Since the minimum mass of Gl 581c and Gl 581d is about 5.0 and 8.0 M_{\oplus} (M_{\oplus} is the Earth mass), respectively, below the boundary of 10 M_{\oplus} between terrestrial and giant gas planets, they are considered to be two terrestrial super-Earth planets. The discovery led to great interest in establishing whether these exoplanets are habitable. von Bloh et al. (2007) and Selsis et al. (2007) estimated whether the two exoplanets are inside the habitable zone of M-dwarf Gl 581 from different points of views. The former study focused on a photosynthesis-sustaining habitable zone that is determined by the limits of photosynthetic life on the planetary surface, and the latter defined the habitable distance using radiative-convective model results. Both suggested that Gl 581c is too close to the M-dwarf and outside the habitable zone, and that Gl 581d is located at the outer edge of the habitable zone and is likely to be habitable.

The habitability of a planet can be determined by various factors, as extensively discussed in the above papers. Among them, a critical and necessary condition is the existence of permanent liquid water on the surface of a planet, which requires that the surface temperature (T_s) is above the freezing point of water (273 K). The temperature T_s of a planet is largely constrained by the radiative properties of atmospheric compositions, in addition to stellar radiation, the distance between the planet and its parent star, surface albedo, and so on. Major greenhouse gases, such as CO₂ and H₂O, can greatly increase T_s by means of their greenhouse effects, as it is well known that the greenhouse effect increases T_s by about 530 K and 33 K for Venus and Earth, respectively. Assuming that the planets Gl 581c and

Gl 581d are terrestrial, and have similar processes in outgassing CO₂ and H₂O as those of the terrestrial planets in the solar system, the two major greenhouse gases would have important influences in determining T_s and thus the habitability of the two exoplanets.

We note that greenhouse gases for the solar system have anti-greenhouse effects for M-dwarf systems because M-dwarfs have much lower effective temperatures. For example, Gl 581 has an effective temperature of about 3200 K (Udry et al. 2007), much lower than the Sun's value of 5800 K. According to Wien's law, the radiation spectrum of Gl 581 peaks at about 0.9 μm in wavelength, which is in the near-infrared (NIR) region. This redshift of the radiation spectrum causes absorption of incident stellar radiation by CO₂ and H₂O in the upper atmospheres of planets orbiting M-dwarfs because both CO₂ and H₂O have strong absorption bands in the NIR region (Goody & Yung 1989; Yung & DeMore 1999). Thus, for M-dwarf stellar radiation, both CO₂ and H₂O have not only greenhouse effects but also anti-greenhouse effects. It is similar to the anti-greenhouse effect of organic aerosols in the atmosphere of Titan (McKay et al. 1991). The latter warms the upper atmosphere, but cools the surface of planets orbiting M-dwarfs. This is unlike the situation in the solar system. Solar radiation has a peak wavelength of about 0.55 μm in the visible region. Both CO₂ and H₂O are nearly transparent to solar radiation and trap only infrared radiation from planets, which warms the planet's surface. Therefore, to assess how surface temperatures of Gl 581c and Gl 581d are constrained by atmospheric radiative processes we need to perform quantitative calculations with realistic radiation transfer models. Wordsworth et al. (2010) showed that less than about 10 bars of CO₂ is sufficient to maintain a global mean T_s of Gl 581d above the freezing point of water with a radiative-convective model,

Table 1. Parameters used in the present study.

Planets	M_{\min}/M_{\oplus}	r/r_{\oplus}	$g(\text{m s}^{-2})$	$\Gamma_d (\text{K km}^{-1})$	$d (\text{AU})$	$S_0 (\text{W m}^{-2})$	$c_p (\text{J K}^{-1} \text{kg}^{-1})$
Gl 581c	5.06	1.8	15.3	15.3	0.073	955.83	1003
Gl 581d	8.30	2.0	20.3	23.9	0.210	103.09	850

Notes. M_{\oplus} is Earth's mass, r_{\oplus} is Earth's radius, M_{\min} and r are planets' minimum mass and radius, respectively. g is surface gravity, Γ_d is the dry adiabatic lapse rate, d is the mean distance between the star and planets, and S_0 is the average stellar flux.

and that the CO_2 threshold is about 30 bars for conservative conditions. In another independent study, von Paris et al. (2010) obtained similar results, but with a slightly lower CO_2 threshold.

In the present paper, we use a different radiative-convective model to investigate the CO_2 threshold of Gl 581d to increase its T_s above the freezing point of water and examine both greenhouse and anti-greenhouse effects of CO_2 and H_2O in the Gliese 581 system. In addition, we also study radiative constraints on the habitability of Gl 581c using another radiative-convective model. In contrast to Gl 581d, Gl 581c is much closer to its parent star. Thus, the key concern for Gl 581c is whether its surface temperature is below the runaway greenhouse threshold (340 K) under conditions of a weak greenhouse effect and high planetary albedo. Models are described in Sect. 2. Simulations results are presented in Sect. 3. Discussion and conclusions are summarized in Sect. 4.

2. Radiative-convective model

Two radiative-convective models are used in the present study. One model is used for simulations for Gl 581d. The model was originally developed by Kasting et al. (1984a,b) for simulating dense planetary atmospheres with high levels of CO_2 , and later modified by Toon et al. (1989), Pavlov et al. (2000), Mischna et al. (2000), and others. The model takes into account the effects of collision-induced absorption by CO_2 , pressure-induced broadening of CO_2 and H_2O absorption, and Rayleigh scattering by CO_2 , and was used in radiation transfer and surface temperature simulations of the early Earth's atmosphere that likely had very high levels of CO_2 (Kasting & Ackerman 1986). It has 38 spectral intervals in the visible and near-infrared radiation and 55 spectral intervals in the thermal-infrared. Infrared absorption by CO_2 and water vapor is calculated from the Air Force Geophysical Laboratory tape. Hereafter, the model is denoted as VPL¹.

The other model is the Santa Barbara DISORT Atmospheric Radiative Transfer (SBDART) model (Ricchiazzi et al. 1998), which is based on low-resolution band models developed for the LOWTRAN 7 atmospheric transmission code. The band models provide clear-sky atmospheric transmission from 0 to 50000 cm^{-1} and are derived from detailed line-by-line calculations with a resolution of 20 cm^{-1} . Convective adjustment is added to the model using the dry adiabatic lapse rate. Since our simulations for Gl 581c have atmospheric conditions similar to that of the Earth atmosphere, SBDART is a suitable model.

The radiation spectrum of Gl 581 is given by the Planck function, assuming that it is a blackbody. On the basis of updated observations (Mayor et al. 2009), we use a distance of 0.21 AU between Gl 581d and its parent star rather than 0.25 AU (Udry et al. 2007). The other parameters used here are all

derived from observations documented in Udry et al. (2007) and Mayor et al. (2009), which are listed in Table 1. In the simulations of Gl 581d below, the volume mixing ratio of CO_2 is set to 96%, similar to that in the atmosphere of Venus, and the surface albedo (A_s) equals 0.15, close to the terrestrial value, except for the specified cases. The zenith angle is set to 60° following Manabe & Strickler (1964). Thus, the calculated surface temperature approximately represents the global mean surface temperature. When determining the dry adiabatic lapse rate, $\Gamma_d = g/c_p$, for Gl 581d, the specific heat capacity c_p is taken to be 850 $\text{J K}^{-1} \text{kg}^{-1}$ because we assume that its atmosphere is CO_2 dominant, and its variations with altitude (pressure and temperature) are neglected (Wordsworth et al. 2010). For Gl 581c, c_p is taken as 1003 $\text{J K}^{-1} \text{kg}^{-1}$ assuming that its atmospheric composition is similar to that of Earth's atmosphere. For the VPL model, 101 vertical levels in pressure coordinates are used, and for the SBDART model 50 vertical levels are used. In all simulations, models are integrated by performing iterations in which time is varied until they reach equilibrium.

3. Simulation results

3.1. Simulations for Gl 581d

Two types of simulations are performed for Gl 581d. The first type of simulations has a dry atmosphere (no water vapor). The simulations indicate how the T_s of Gl 581d changes with increasing CO_2 and how the anti-greenhouse effect of CO_2 can be tested. The second type of simulations incorporates a moist atmosphere. Various levels of water-vapor mixing ratios are added to the model to test both the greenhouse and anti-greenhouse effects of water vapor on T_s . We also perform simulations with fixed relative humidity (RH), which more realistically show the dependence of water-vapor concentration on CO_2 levels and their greenhouse effects on T_s .

Figure 1a shows simulated vertical temperature profiles with surface air pressure (P_s) ranging from 1 to 50 bars, that is, CO_2 partial pressure varies from 0.96 to 48 bars. It is found that T_s increases from 217.2 K to 375.6 K with increasing CO_2 . For $P_s = 5$ bar, T_s is 257.8 K. For $P_s = 10$ bar, T_s is 290.8 K, indicating that less than 10 bars of CO_2 is sufficient to maintain T_s of Gl 581d above the freezing point of water. Linear interpolation yields a value of $P_s = 7.32$ bars for $T_s = 273$ K, which is equivalent to about 7.0 bars of CO_2 , when considering the 96% CO_2 mixing ratio.

One problem associated with the results in Fig. 1a is that CO_2 condensation is not considered. From Fig. 1a, one can find that vertical temperature profiles fall into the solid-phase region of CO_2 between about 0.2 and 7.0 bars, indicating that there would be a CO_2 condensation in this layer. CO_2 condensation releases latent heat, which warms the layer where condensation forms and alters the vertical thermal structures and surface temperatures. Simulations in which CO_2 condensation is considered indeed contain a warmed layer between 0.2 and 7.0 bars

¹ Detailed descriptions of the model and relevant references can be found at

<http://vpl.astro.washington.edu/sci/AntiModels/models09.html>




Fig. 1. Simulated vertical temperature profiles for a dry atmosphere of Gl 581d with different CO₂ levels. Dashed-dotted lines indicate CO₂ phase diagram. **a)** CO₂ condensation is not considered, and **b)** CO₂ condensation is considered.

and exhibit decreases in T_s (Fig. 1b). Figure 1b displays lower surface temperatures than Fig. 1a. The reason why surface temperatures decrease is because the warmed layer due to CO₂ condensation emits more outgoing infrared radiation at higher temperatures ($F = \sigma T^4$, where σ is the Stefan-Boltzmann constant and T is air temperature), causing a weakened greenhouse effect on the surface. As a result, the CO₂ threshold is higher than that without considering CO₂ condensation. One can find from Fig. 1b that about 9.6 bars of CO₂, i.e., $P_s = 10$ bar, is required for T_s above 273 K.

The anti-greenhouse effect of CO₂ is illustrated in Fig. 2. The black line in Fig. 2a shows the radiation spectrum of M-dwarf Gl 581 at the top of the Gl 581d atmosphere. It has a peak wavelength at about 0.9 μm . The red line is the incident radiation spectrum on the surface of Gl 581d for $P_s = 5$ bar. The red line clearly shows that a significant portion of incident stellar radiation with wavelength greater than 1 μm is absorbed by CO₂ before it reaches the surface. The anti-greenhouse effect of CO₂ can be measured by the percentage of incident radiation absorbed by the atmosphere. The absorption percentage as a function of P_s is shown in Fig. 2b. For $P_s = 1$ bar ($P_{\text{CO}_2} = 0.96$ bar), the absorption percentage is about 20%. For $P_s = 100$ bar ($P_{\text{CO}_2} = 96$ bar), the absorption percentage is up to 46%. Because of the anti-greenhouse effect, the upper atmosphere is warmed. The inversion layers between 0.6 and 2 bars in Fig. 1a and between 0.6 and 4 bars in Fig. 1b are all due to the anti-greenhouse effect.

Rayleigh scattering of CO₂ plays an important role in reducing incident radiation for atmospheres with high-level CO₂, as shown by Kasting and Ackerman (1986) for the early Earth atmosphere. For M-dwarf stellar radiation, Rayleigh scattering




Fig. 2. **a)** Incident radiative spectrum of M-dwarf Gliese 581 at the top of the atmosphere (black line) and on the surface of Gl 581d for the dry condition with $P_s = 5$ bar (CO₂ condensation is not taken into account). The red and blue lines indicate surface incident radiative spectra with and without Rayleigh scattering, respectively. **b)** Red line: percentage of incident radiation absorbed by CO₂ as a function of P_s , and blue line: percentage of incident radiation absorbed and scattered by CO₂.

of CO₂ is weaker because of the red shift of stellar radiation. Nonetheless, Rayleigh scattering of CO₂ still has important effects in the visible region, especially for high levels of CO₂. In Fig. 2a, the blue line shows the spectrum of incident radiation on the surface as CO₂ absorption and Rayleigh scattering are all considered. One can differentiate between blue and red lines in the region with wavelengths shorter than 1.25 μm . For longer wavelengths, Rayleigh scattering quickly decays because its effect is proportional to λ^{-4} , where λ is the wavelength of incident radiation. The blue line in Fig. 2b shows the percentage of incident radiation absorbed and scattered by CO₂ as a function P_s . Comparison of the blue line with the red line reveals that the effect of Rayleigh scattering becomes more and more important as CO₂ increases. For $P_s = 1$ bar, Rayleigh scattering has very weak effect. As P_s increases to 10 and 50 bars, it leads to a reduction in the incident radiation by 10% and 20%, respectively.

The above results are for a dry atmosphere. Assuming that Gl 581d has similar outgassing processes to that of terrestrial planets in the solar system, its atmosphere should also include water vapor. To examine how the greenhouse effect of water vapor, and that of CO₂, influences T_s , we consider $P_s = 5$ bar and fix the surface water-vapor mixing ratios to 1, 10, 100, and 1000 ppmv, respectively. Water-vapor mixing ratios decay exponentially with altitudes to the same value of 1 ppmv at 0.5 bar and remain constant above this pressure level, as shown in Fig. 3b. In these simulations, convective adjustment is used with the dry adiabatic lapse rate, $\Gamma_d = g/c_p = 23.88 \text{ K km}^{-1}$. Simulation results are shown in Fig. 3a. The greenhouse effect of water vapor is strong. Even 1 ppmv of water vapor can increase T_s from 257.8 K for the dry atmosphere (see the green line in




Fig. 3. **a)** Simulated vertical temperature profiles for a moist atmosphere of Gl 581d, with $P_s = 5$ bar and a dry adiabatic lapse rate of 23.88 K km^{-1} . CO_2 and water vapor condensation is not considered. Dotted-line indicates the water-vapor saturation line, and the dashed and dotted-line indicates the CO_2 phase-diagram between gas and solid phases. **b)** Vertical distributions of water-vapor mixing ratios.

Fig. 1a) to 263.3 K. For 10, 100, and 1000 ppmv of water vapor, the corresponding T_s is 263.4, 268.3, and 274.2 K, respectively. The results suggest that a lower CO_2 threshold is required as water vapor is considered.

The anti-greenhouse effect also becomes much stronger when water vapor is included. A comparison of Fig. 4a with Fig. 2a indicates that more incident radiation is absorbed by the atmosphere, especially in the region from the peak wavelength to the near-infrared region. From Fig. 4b, one can infer that the absorption percentage is about 32% as 1 ppmv of water vapor is added to the model, which is about 1% higher than the dry case for $P_s = 5$ bar (see Fig. 2b). For 1000 ppmv of water vapor, about 44% of the incident radiation is absorbed by the atmosphere, which represents an increase of about 13%.

We note that water vapor concentration is constrained by ambient air temperatures, that is, water vapor condenses when it is saturated. Thus, one cannot arbitrarily add water vapor to the model. Figure 3a already shows super-saturation, in which temperature profiles are below the saturation line at some levels. It suggests that the results in Fig. 3a may not be realistic. A more sophisticated way of examining how T_s is determined by water vapor, as well as CO_2 , is to use the method of fixed relative humidity (RH). In this way, water vapor concentration varies with air temperatures for different CO_2 levels. In these simulations below, the vertical distribution of RH in the troposphere is given by the formula of Manabe and Wetherald (1967) for Earth's atmosphere. This method was used to study Earth's early atmosphere by Kasting and Ackerman (1986).

Simulations with fixed RH are performed with three values of surface albedo, i.e., $A_s = 0.15, 0.20,$ and 0.30 . In these simulations, moist adiabatic processes are applied, and CO_2 condensation is also considered. Figure 5a shows vertical temperature profiles for $A_s = 0.15$. Comparison of Fig. 5a with Fig. 1b shows that the surface temperature T_s is much higher when water vapor is included in the model. For example, for $P_s = 10$ bar, T_s is 276.7 K for the dry atmosphere, while it is 291.9 K for the moist atmosphere. This difference is due to the greenhouse effect of water vapor. However, the surface temperature for $P_s = 5$ bar, 257.9 K, is lower than in Fig. 3a because of the condensation of CO_2 and water vapor, as well as lower lapse rates in the moist




Fig. 4. **a)** Same as Fig. 2a, except for that 1000 ppmv of water vapor is added to the model. **b)** Percentage of absorption and absorption with Rayleigh scattering as a function of water vapor mixing ratio.

atmosphere. As mentioned before, the condensation of CO_2 and water vapor warms the upper troposphere, causing a weakened greenhouse effect. The temperature lapse rate for moist adiabatic processes is usually lower than that in the case of dry adiabatic processes. A lower lapse rate causes a weaker greenhouse effect because outgoing infrared radiation from the upper troposphere is stronger when the air temperature is higher than for a dry adiabatic lapse rate, leading to a decrease in T_s .

The temperature T_s as a function of P_s , under conditions of CO_2 condensation, moist adiabatic processes, and fixed RH, is plotted in Fig. 5b. The solid-square line in Fig. 5b is for $A_s = 0.15$. It crosses the dotted-line ($T_s = 273 \text{ K}$) at about $P_s = 7.0$ bar, indicating that a minimum CO_2 level of 6.7 bars is required for T_s to be maintained above the freezing point of water. To reach the Earth's global-mean surface temperature today (288 K), CO_2 has to be about 8.64 bars ($P_s = 9.0$ bar). For $P_s = 50$ bar, T_s is 362.9 K. One might worry that Gl 581d would undergo runaway greenhouse for $P_s = 50$ bar since the corresponding T_s is above the threshold of 340 K (Ingersoll 1969). However, 340 K is the threshold for 1 bar of surface air pressure, and the threshold increases with P_s , as pointed out by Selsis et al. (2007). For surface albedos of 0.20 and 0.30, P_s corresponding to 273 K is about 8.0 and 12.0 bars, respectively, equivalent to about 7.7 and 11.5 bars of CO_2 . We note that the three lines tend to converge for $P_s = 50$ bar, which indicates that the effect of surface albedo on T_s becomes less important for high CO_2 levels. This is because incident stellar radiation is mainly absorbed and scattered by the atmosphere (CO_2 and water vapor) for high levels of CO_2 , and a very minor part of incident radiation can reach the surface. Thus, T_s is mainly determined by the greenhouse effect of the atmosphere, rather than by incident radiation reaching the surface. The issue of surface albedo effect at very high levels of CO_2 was also studied by Selsis et al. (2007).




Fig. 5. **a)** Simulated vertical temperature profiles for a moist atmosphere for Gl 581d with fixed RH and $A_s = 0.15$. **b)** Simulated T_s as a function of P_s with fixed RH and moist adiabatic lapse rates.

Wordsworth et al. (2010) and von Paris et al. (2010) independently studied the CO_2 threshold for Gl 581d (Hereafter, the two paper are referred as WW and vP, respectively). WW used a different radiative-convective model from ours, while vP used the same model as ours, but with different parameterizations of absorption for infrared radiation. Here, we compare these results and assess the cause of differences among these models. Figs. 6a–c compare our results and WW’s for dry, moist, and water-vapor saturation atmospheres, respectively. To provide the comparison, simulations are carried out using the same parameters as that in WW and vP, such as surface albedo, temperature lapse rate, the vertical distribution of relative humidity, and so on. Figure 6a shows the comparison for the dry case (pure CO_2). As P_s is lower than 2.5 bars, T_s in the VPL model is slightly lower than that in WW for the same level of CO_2 . For example, the difference in T_s is about 3 K for $P_s = 1.0$ bar. However, as P_s is greater than 2.5 bars, T_s in the VPL model is higher than in WW, and the difference in T_s between the two models increases with increasing P_s . For $P_s = 20$ bars, T_s in the VPL model is about 6 K higher than in WW (about 292 versus 286 K). For low levels of CO_2 , the WW model appears to have a slightly stronger greenhouse effect than the VPL model probably because Wordsworth et al. (2010) used the HITRAN 2008 database for CO_2 absorption, which differs from that in the VPL model. For high levels of CO_2 , collision-induced absorption of CO_2 becomes important and causes an enhanced greenhouse effect for CO_2 . The comparison indicates that the parameterization for collision-induced absorption of CO_2 in WW, which is based on new measurement results (Baranov et al. 2004; Gruszka & Borysow 1998), tends to reduce greenhouse

warming for high levels of CO_2 . For the moist case (Fig. 6b), T_s in the VPL model is systematically lower than that in WW, and the difference also becomes greater with increasing P_s . For $P_s = 20$ bars, the T_s in the VPL model is about 13 K lower than that in WW, about 332 K versus 345 K. This suggests that water vapor has a stronger greenhouse effect in WW’s model than in the VPL model, which increases T_s in WW’s model from 6 K lower than in the VPL model in the dry case to 13 K higher than in the VPL model in the moist case for $P_s = 20$ bars. For the saturation case (Fig. 6c), T_s in WW is much higher than in the VPL model. For $P_s = 20$ bars, T_s in WW is about 19 K higher than in the VPL model, about 367 K versus 348 K. This is again indicative of a stronger greenhouse effect of water vapor in WW’s model than in the VPL model.

Although vP used the same model as ours, i.e., the VPL model, they changed both the parameter values and parameterizations of the model. In particular, they recomputed k -correlated coefficients with their own line-by-line model in order to achieve a wider coverage of pressure and temperature. A comparison of our results with those of vP is shown in Fig. 6d. Our results (crosses) are also systematically lower than vP’s, and the difference in T_s ranges from about 13 K for $P_s = 1$ bar to about 30 K for $P_s = 5$ or 10 bars. The new parameterization of water vapor absorption may play an important role in causing higher temperatures in vP than in the VPL model, especially for high levels of CO_2 . However, it does not appear to be the major reason for causing a difference in T_s for low levels of CO_2 , at which T_s is so low that little water vapor in the atmosphere can be present. Thus, other changes in parameters and parameterizations for CO_2 are also responsible for the stronger greenhouse effect in vP. In Fig. 6d, we also determine the difference between cases of CO_2 super-saturation and condensation. It is found that the CO_2 super-saturation causes about a 7 K increase in T_s for $P_s = 20$ bars. It is difficult to directly compare the results between WW and vP because they used very different parameters. One major reason that causes higher T_s in vP than in WW is probably because surface albedo in vP is much lower than in WW (0.13 versus 0.2). For a moist atmosphere, the models used overall by WW and vP have stronger greenhouse effects than the VPL model.

3.2. Simulations for Gl 581c

Gl 581c is much closer to its parent star than Gl 581d, at a distance of about 0.073 AU. It receives 30% more incident stellar radiation than Venus today (Udry et al. 2007; Selsis et al. 2007). Because the radiative equilibrium temperature of Gl 581c is already as high as 300 K for an albedo of 0.5 (Udry et al. 2007), the greenhouse effect can readily increase its T_s above the runaway greenhouse threshold, suggesting that a high planetary albedo and weak greenhouse effect are necessary for Gl 581c to be habitable. The following simulations test whether these two conditions can be realized.

First, we test the condition of weak greenhouse effect by using a low CO_2 level in the model. The dash-dotted line in Fig. 7 is the simulated temperature profile for a dry atmosphere with conditions of $P_s = 1$ bar (same as that of Earth), 50 ppmv of CO_2 , and $A_s = 0.5$. The corresponding T_s is 310 K. Second, we add water vapor to the model. The vertical distribution of a water-vapor mixing ratio is shown in Fig. 7b (dashed-line), where the surface water-vapor mixing ratio is the same as that in Earth’s atmosphere (US standard atmosphere, 1976). The reason why the water-vapor mixing ratio is set to be uniform above 700 hPa is because the convective layer is very thin and can only reach




Fig. 6. Comparison of our simulation results for Gl 581d with that in WW and vP. Plots a-c compare the results of WW with our results, and plot d compares the result of vP with our results. **a)** Dry atmosphere with considering CO₂ condensation, **b)** moist atmosphere with fixed RH, dry adiabatic lapse rate and considering condensation of CO₂, and **c)** same as **b)**, except for saturated water vapor (RH = 100%) in the troposphere. In plots **a)**–**c)**, $A_s = 0.20$, gravity $g = 20 \text{ m s}^{-2}$. The dry atmosphere here is pure CO₂, and the moist atmosphere is dominated by CO₂ and contains a minor component of water vapor. The vertical distribution of RH is the Manabe-Wetherald type, with surface RH = 0.77. In plot **d)**, parameters are same as that in vP, that is, $A_s = 0.13$, gravity $g = 23.76 \text{ m s}^{-2}$, CO₂ concentration is 95%, the vertical distribution of RH is the Manabe-Wetherald type with surface RH = 0.80, and CO₂ is super-saturated. Here, S_{sat} indicates super-saturation of CO₂. In all the plots, T_s values by WW and vP are estimated from their figures.

700 hPa. When this amount of water vapor is added, T_s rapidly increases to 331 K (dashed-line in Fig. 7a). Although it remains below 340 K, the surface RH is too low, about 3.4%, to allow the existence of liquid water on the surface. To maintain permanent liquid water on Gl 581c, the surface RH must be comparable to that of Earth. However, simulations in which the fixed RH is almost identical to that used in Fig. 5 show that T_s rises quickly above 340 K, and the model cannot reach equilibrium, which is indicative of runaway greenhouse. The above results suggest that the condition of a weak greenhouse effect can hardly be realized.

There are two possibilities of a high planetary albedo for Gl 581c: a large coverage of either snow-ice or thick clouds (Selsis et al. 2007). The former is unlikely because CO₂ accumulation due to volcanic eruptions would eventually melt the snow and ice, resulting in a runaway greenhouse, even if the planet were cold initially. A high albedo due to thick clouds might be more likely because a water-rich, warm planet would have deep convective clouds (Selsis et al. 2007). However, simulations indicate that the assumption may also be unrealistic because the

cloud optical thickness has to be as high as 270 to keep T_s below the threshold of runaway greenhouse.

The solid line in Fig. 7 is the simulated temperature profile, where the cloud optical thickness is 270, the effective radius of cloud drops is 20 μm (similar to terrestrial clouds), and RH is fixed as above. The simulation yields a T_s of about 338 K, very close to the threshold of runaway greenhouse. Such a high optical thickness corresponds to a very dark world, with which there is not only little direct incident radiation but also very little scattered shortwave radiation that can reach the surface. Our calculations indeed show that only 8.76 W m^{-2} of scattered radiation can reach the surface (total incident radiation is about 955.83 W m^{-2}), while direct incident radiation is negligible. The problem is that such thick clouds as these may not be realistic. Moreover, the above simulation is based on the assumption of 100% cloud coverage. A more realistic possibility is that strong convection must be accompanied by descending regions where there are few or no clouds. Thus, an even higher optical thickness of clouds is needed to satisfy the requirement of a high planetary albedo.




Fig. 7. a) Simulated vertical temperature profiles for Gl 581c. Dash and dotted-line: dry atmosphere with $P_s = 1$ bar and 50 ppmv of CO_2 , Dashed-line: moist atmosphere with $P_s = 1$ bar, 50 ppmv of CO_2 , and same water-vapor mixing ratio as in the Earth atmosphere, and solid line: moist atmosphere with $P_s = 1$ bar, 50 ppmv CO_2 , fixed relative humidity same as in Manabe and Wetherald (1967). A cloud layer is located between 700 and 100 hPa marked by horizontal dotted-lines. b) Vertical profiles of water-vapor volume mixing ratio. The solid and dashed-lines in b) correspond to the solid and dashed-lines in a), respectively.

The lack of an effective cold trap for water vapor is also a problem. The dashed and solid lines in Fig. 7 all indicate that the lowest temperatures are higher than 273 K in the presence of water vapor, suggesting a reduction in the effectiveness of the cold trap for water. It would allow more water vapor into the stratosphere relative to that crossing the tropopause of Earth's atmosphere because there is no dehydration process such as that near Earth's tropopause. As a result, water loss is much faster due to the stronger photolysis of water vapor in the middle atmosphere (Kasting et al. 1993). In addition, Gl 581c may have a much denser atmosphere and higher levels of greenhouse gases since it has a much higher mass than Earth. If it is the case, Gl 581c must have experienced runaway greenhouse, just like Venus.

4. Discussion and conclusions

Using a real-gas radiative-convective model, we have studied the CO_2 threshold for maintaining the surface temperature of Gl 581d above the freezing point of water. For a dry atmosphere affected by CO_2 condensation and $A_s = 0.15$, our simulations demonstrate that at least 9.6 bars of CO_2 are required. For a moist atmosphere with fixed RH and $A_s = 0.15$, the CO_2 threshold is lowered to 6.7 bars because of the greenhouse effect of water vapor. For $A_s = 0.30$ and a moist atmosphere with fixed RH, which is close to Earth's planetary albedo, the CO_2 threshold is about 11.5 bars. The CO_2 threshold in our simulations is slightly higher than that in Wordsworth et al. (2010) and von Paris et al. (2010) because their models all have a stronger greenhouse effect for a moist atmosphere. Our comparison indicates that one of the major factors in causing the differences in T_s is presumably the new parameterizations for CO_2 and water vapor absorption in their models. We have also found that collision-induced CO_2 absorption in Wordsworth et al. (2010) has slightly weaker greenhouse effect.

In our simulations for Gl 581d, we have not considered the radiative effect of clouds of both CO_2 and water, which also play important roles in constraining T_s . Liquid water clouds reflect incident shortwave radiation and cool the surface, whereas ice clouds of CO_2 and water have greenhouse effects and warm the surface. In particular, CO_2 ice clouds in a high- CO_2 atmosphere

have a strong greenhouse effect on the surface. As shown by Forget and Pierrehumbert (1997), CO_2 ice clouds were a possible factor in causing the greenhouse effect and warming early Mars' surface. We do not study the influence of clouds on the T_s of Gl 581d because the way in which clouds vary depending on CO_2 levels is not known for the one-dimensional model (Kasting & Ackerman 1986).

It is not unrealistic for Gl 581d to contain CO_2 at around 10 bars in its atmosphere because it has a minimum mass of about 10 times greater than that of Venus, which has up to 90 bars of CO_2 in its atmosphere. However, the maintenance of the high level of CO_2 in the atmosphere of Venus is due to the lack of liquid water, which allows CO_2 , produced in turn by volcano eruptions, to become accumulated in the atmosphere. As long as there is permanent liquid water on the surface of a planet, the water cycle becomes active, removing CO_2 from the atmosphere by means of weathering reactions and leading to a low level of CO_2 (Walker et al. 1981; Kasting & Ackerman 1986; Kasting 1988). For example, the formation of the Snowball Earth in the Neoproterozoic era was proposed to be caused when CO_2 dropped to about 100 ppmv in Earth's atmosphere because of active weathering reactions (Hoffman & Schrag 2000; Pierrehumbert 2004). The question then arises of how CO_2 can be maintained at a level around 10 bars for Gl 581d in the presence of liquid water. One possibility is that the surface of Gl 581d is mostly covered by ocean, in which case the carbonate-silicate cycle could be inefficient, as suggested by Selsis et al. (2007). It is likely that Gl 581d is mostly covered by ocean because as the mass of the planet increases so does the water-to-surface ratio (Lissauer 1999). Given that Gl 581d resembles an aqua-planet, its surface albedo would be correspondingly lower, and the required minimum CO_2 level would also be lower than the above results. It is also possible that the climate system of Gl 581d oscillates between the snowball and habitable regimes driven by the carbonate-silicate cycle.

We have found that our simulations imply that Gl 581c is too hot to be habitable. Since its radiative equilibrium temperature is already very high, even a weak greenhouse effect can readily increase its surface temperature above the threshold of runaway greenhouse. The high planetary albedo caused by extremely thick clouds seems unrealistic. The lack of an effective

cold trap for water would lead to rapid loss of water. All of these results suggest that Gl 581c had very likely undergone runaway greenhouse like Venus. We note that all our simulations have been performed without considering either the eccentricity or the obliquity. The results here also do not reject the possibility of habitable areas in the transition regions between day and night hemispheres, where the temperature is neither too high nor too low, given that Gl 581c is a synchronously rotating planet. More detailed and qualitative discussion about these aspects can be found in Selsis et al. (2007).

The model predictions discussed in this paper are testable by observations. Given that Gl 581c is a Venus-like planet because it has experienced runaway greenhouse, it should display the distinctive characteristics of Venus, such as a high albedo with blue and ultraviolet broadband absorption, SO₂ absorption band due to H₂SO₄ clouds, and the absence of H₂O absorption bands, which are detectable (Mills et al. 2007). The presence or absence of H₂O and CO₂ on Gliese 581c and Gl 581d may be established by observations such as those for the hot Jupiter HD 189733b (Tinetti et al. 2007; Swain et al. 2009).

Acknowledgements. We are grateful to Prof. Douglas Lin who made helpful comments on an early version of the paper. We thank the anonymous reviewer for careful reviews and very helpful comments, which lead to important corrections and improvements of the paper. This work is supported by the National Basic Research Program of China (973 Program, 2010CB428606), the National Natural Science Foundation of China (40875042, 41025018), and the Ministry of Education of China (20070001002).

References

- Baranov, Y. I., Lafferty, W. J., & Fraser, G. T. 2004, *J. Mol. Spectrosc.*, 228, 432
- Forget, F., & Pierrehumbert, R. T. 1997, *Science*, 278, 1273
- Goody, R. M., & Yung, Y. L. 1989, *Atmospheric Radiation: Theoretical Basis*. 2nd Ed. (Oxford University Press)
- Gruszka, M. & Borysow, A. 1998, *Molecul. Phys.*, 93, 1007
- Hoffman, P. F., & Schrag, D. P. 2000, *Sci. Am.*, 282, 62
- Ingersoll, P. 1969, *J. Atmos. Sci.*, 26, 1191
- Kasting, J. F. 1988, *Icarus*, 74, 472
- Kasting, J. F., & Ackerman, T. P. 1986, *Science*, 234, 1383
- Kasting, J. F., Pollack, J. B., & Ackerman, T. P. 1984a, *Icarus*, 57, 335
- Kasting, J. F., Pollack, J. B., & Crisp, D. 1984b, *J. Atmos. Chem.*, 1, 403
- Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, *Icarus*, 101, 108
- Li, K.-F., Pahlevan, K., Kirschvink, J. L., & Yung, Y. L. 2009, *Proc. Nat. Acad. Sci., USA*, 106, 9576
- Lissauer, J. J. 1999, *Nature*, 402, C11
- Manabe, S., & Strickler, R. F. 1964, *J. Atmos. Sci.*, 21, 361
- Manabe, S., & Wetherald, R. T. 1967, *J. Atmos. Sci.*, 24, 241
- Mayor, M., Bonfils, X., Forveille, T., et al. 2009, *A&A*, 507, 487
- McKay, C. P., Pollack, J. B., & Courtin, R. 1991, *Science*, 253, 1118
- Mills, F. P., Esposito, L. W., & Yung, Y. L. 2007, in *Exploring Venus as a Terrestrial Planet*, ed. L. W., Esposito, E., Stofan, T., Cravens, American Geophysical Union, Washington, DC, 73
- Mischna, M. A., Pavlov, A. A., & Kasting, J. F. 2000, *Icarus*, 145, 546
- Pavlov, A. A., Kasting, J. F., Brown, L. L., Rages, K. A., & Freedman, R. 2000, *J. Geophys. Res.*, 105, 11981
- Pierrehumbert, R. T. 2004, *Nature*, 429, 646
- Ricchiazzi, P., Yang, S., Gautier, C., & Sowle, D. 1998, 79, 2101
- Selsis, F., Kasting, J. F., Levrard, B., et al. 2007, *A&A*, 476, 1373
- Swain, M. R., Vasisht, G., Tinetti, G., et al. 2009, *Astrophys. J.*, 690, L114
- Tinetti, G., Vidal-Madjar, A., Liang, M. C., et al. 2007, *Nature*, 448, 169
- Toon, O. B., McKay, C. P., Ackerman, T. P., & Santhanam, K. 1989, *J. Geophys. Res.*, 94, 16287-16301
- Udry, S., Bonfils, X., Delfosse, X., et al. 2007, *A&A*, 469, L43
- US standard atmosphere, 1976, US Government printing office, Washington DC
- Walker, J. C. G., Hays, P. B., & Kasting, J. F. 1981, *J. Geophys. Res.*, 86, 9776
- von Bloh, W., Bounama, C., Cuntz, M., & Franck, S. 2007, *A&A*, 476, 1365
- von Paris, P., Gebauer, S., Godolt, M., et al. 2010, accepted by *A&A*, 522, A23
- Wordsworth, R. D., Forget, F., Selsis, F., et al. 2010, *A&A*, in press
- Yung, Y. L., & DeMore, W. D. 1999, *Photochemistry of Planetary Atmospheres* (Oxford University Press)